

Datasheet for ABIN6719997

IL-2 ELISA Kit[Go to Product page](#)**1** Image

Overview

Quantity:	1 kit
Target:	IL-2 (IL2)
Reactivity:	Rat
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

Product Details

Purpose:	Rat IL-2 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin - citrate), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant rat IL-2. There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 5 pg/mL
Components:	<ul style="list-style-type: none">• Antibody-coated 96-well plate• Target Protein Standard• Detection antibody

Product Details

- Detection reagent
- Diluent buffers
- Wash buffers
- Substrate Solution
- Stop solutions
- Adhesive covers

Target Details

Target: IL-2 (IL2)

Alternative Name: IL-2 ([IL2 Products](#))

Background: Synonyms: Aldesleukin, IL 2, IL-2, IL2, IL2_HUMAN, Interleukin 2, Interleukin-2, Interleukin2, Involved in regulation of T cell clonal expansion, Lymphokine, T cell growth factor, T-cell growth factor, TCGF

Background: Interleukin-2(IL2), formerly referred to as T-cell growth factor, is a powerfully immunoregulatory lymphokine that is produced by lectin- or antigen-activated T cells. It is produced not only by mature T lymphocytes on stimulation but also constitutively by certain T-cell lymphoma cell lines. The lymphokine interleukin-2(IL-2) is responsible for autocrine cell cycle progression and regulation of immune responses. IL-2 expression in mature thymocytes and T cells has been found to be tightly controlled by monoallelic expression. IL-2 can act as a growth hormone for both B and T lymphocytes. The human gene for interleukin 2(IL2) is assigned to chromosome 4.

Gene ID: 116562

NCBI Accession: [NP_446288](#)

UniProt: [P17108](#)

Pathways: [JAK-STAT Signaling](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#), [Activated T Cell Proliferation](#)

Application Details

Application Notes: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL rat IL-2 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of rat cell culture

Application Details

supernates, serum or plasma(heparin, EDTA, citrate) to each empty well. We recommend that each rat IL-2 standard solution and each sample is measured in duplicate.

Comment: Standard: Expression system for standard: E.coli, Immunogen sequence: A21-Q155

Sample Volume: 100 μ L

Plate: Pre-coated

Restrictions: For Research Use only


Handling

Storage: 4 °C,-20 °C

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiry Date: 12 months

Images



ELISA

Image 1. Rat IL-2 Accusignal ELISA Kit Rat IL-2 AccuSignal ELISA Kit standard curve. Assay Range: 31.2pg/ml-2000pg/ml. Sensitivity: <5pg/ml.