

Datasheet for ABIN6720008

**N-Cadherin ELISA Kit****1** Image[Go to Product page](#)

## Overview

Quantity:	1 kit
Target:	N-Cadherin (CDH2)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.78 ng/mL - 50 ng/mL
Minimum Detection Limit:	0.78 ng/mL
Application:	ELISA

## Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric

Characteristics:	<p>Synonyms: CDw325, CDHN, Neural cadherin, N-cadherin, CD325</p> <p>Background: Cadherin-2(CDH2), also known as neural cadherin(NCAD), is a protein that in humans is encoded by the CDH2 gene. It is a classical cadherin from the cadherin superfamily. This gene is mapped to 18q12.1. Cadherin-2 is expressed in the brain, skeletal and cardiac muscle. Cadherin-2 is commonly found in cancer cells and provides a mechanism for transendothelial migration. It is a calcium dependent cell-cell adhesion glycoprotein comprising five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. The protein functions during gastrulation and is required for establishment of</p>
------------------	--

## Product Details

left-right asymmetry. At certain central nervous system synapses, presynaptic to postsynaptic adhesion is mediated at least in part by this gene product.

Gene Name: CDH2

Production: Natural and recombinant human Cadherin-2. There is no detectable cross-reactivity with other relevant proteins.\

Standard: Expression system for standard: NSO, Immunogen sequence: D160-A724

## Target Details

Target:	N-Cadherin (CDH2)
Alternative Name:	Cadherin-2 - N-Cadherin ( <a href="#">CDH2 Products</a> )
Gene ID:	1000
NCBI Accession:	<a href="#">NP_001295105</a>
UniProt:	<a href="#">P19022</a>
Pathways:	<a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Cell-Cell Junction Organization</a> , <a href="#">Synaptic Membrane</a>

## Application Details

Application Notes:	<p>Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 50 ng/mL, 25 ng/mL, 12.5 ng/mL, 6.25 ng/mL, 3.12 ng/mL, 1.56 ng/mL, 0.78 ng/mL human Cadherin-2 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. We recommend that each human Cadherin-2 standard solution and each sample is measured in duplicate.</p> <p>Blood Product Anticoagulant: Heparin Sodium</p> <p>ELISA Dilution: 0.78 ng/mL-50 ng/mL</p>
Sample Volume:	100 µL
Plate:	Pre-coated
Restrictions:	For Research Use only

## Handling

Storage:	RT,4 °C,-20 °C
----------	----------------

Handling

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images

