antibodies -online.com





IGFBP2 ELISA Kit





Overview

Quantity:	1 kit
Target:	IGFBP2
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	125 pg/mL - 8000 pg/mL
Minimum Detection Limit:	125 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen	
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Synonyms: BP 2, BP2, IBP 2, IBP-2, IBP2, IBP2_HUMAN, IGF binding protein 2, IGF BP53, IGF-	
	binding protein 2, IGF-BP53, IGFBP-2, IGFBP2, IGFBP53, Insulin Like Growth Factor Binding	
	Protein 2, Insulin like growth factor binding protein 2 36 kDa, Insulin like growth factor-binding	
	protein 2 precursor, Insulin-like growth factor-binding protein 2	
	Background: Insulin-like growth factor-binding protein 2(IGFBP2), also called IBP2, is a protein	
	that in humans is encoded by the IGFBP2 gene. It is mapped to 2q35. By structural analysis, it	
	showed that the IGFBP2 gene consists of 4 exons with 3 introns of lengths 27.0, 1.0, and 1.9 kb.	

IGFBP2 secreted by metastatic cells recruits endothelia by modulating IGF1 mediated

activation of the IGF type-I receptor on endothelial cells. The IGFBP2/IGF1/IGF1R and

GAS6/MERTK signaling pathways are regulators of cancer-mediated endothelial recruitment.

Gene Name: IGFBP2

Production: Natural and recombinant mouse IGFBP2. There is no detectable cross-reactivity

with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: E35-Q305

Target Details

Target:	IGFBP2
Alternative Name:	IGFBP2 (IGFBP2 Products)
Gene ID:	16008
NCBI Accession:	NP_032368
UniProt:	P47877
Pathways:	Myometrial Relaxation and Contraction, Growth Factor Binding, Activated T Cell Proliferation

Application Details

Application	Notes:

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 8000pg/mL, 4000pg/mL, 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL mouse IGFBP2 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each mouse IGFBP2 standard solution and each sample be measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 125pg/mL-8000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only

Handling


Storage: RT,4 °C,-20 °C

Handling

Storage Comment:

Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Mouse IGFBP2 Accusignal ELISA Kit Mouse IGFBP2 AccuSignal ELISA Kit standard curve. Assay Range: 125pg/ml-8000pg/ml. Sensitivity: <10pg/ml.