

Datasheet for ABIN6720053

IL-4 ELISA Kit**1** Image[Go to Product page](#)

Overview

Quantity:	1 kit
Target:	IL-4 (IL4)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	15.6 pg/mL - 1000 pg/mL
Minimum Detection Limit:	15.6 pg/mL
Application:	ELISA

Product Details

Purpose:	Human IL-4 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant human IL-4 . There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 1.5 pg/mL
Components:	<ul style="list-style-type: none">• Antibody-coated 96-well plate• Target Protein Standard• Detection antibody

Product Details

- Detection reagent
- Diluent buffers
- Wash buffers
- Substrate Solution
- Stop solutions
- Adhesive covers

Target Details

Target:	IL-4 (IL4)
Alternative Name:	IL-4 (IL4 Products)
Background:	<p>Synonyms: B cell growth factor 1, B cell stimulatory factor 1, BCGF 1, BCGF1, Binetrakin, BSF 1, BSF1, IL 4, Interleukin 4, Lymphocyte stimulatory factor 1, MGC79402, Pitrakinra</p> <p>Background: Interleukin-4(IL-4), also known as a B-cell stimulatory factor1(BSF1), is an immunomodulatory cytokine, which can inhibit the growth of tumour cells. The human cDNA contains a single open reading frame encoding a protein of 153 amino acids, including a putative signal peptide. IL-4 may act as an autocrine growth factor in pancreatic cancer cells and also give rise to the possibility that cancer-derived IL-4 may suppress cancer-directed immunosurveillance in vivo in addition to its growth-promoting effects, thereby facilitating pancreatic tumor growth and metastasis. The mouse and human genes and their protein products show structural and functional similarities. The human IL-4 gene, which occurs as a single copy in the haploid genome, is mapped on chromosome 5. The standard product used in this kit is recombinant human IL-4, consisting of 130 amino acids with the molecular mass of 14KDa.</p>
Gene ID:	3565
NCBI Accession:	NP_000580
UniProt:	D4HNR6
Pathways:	JAK-STAT Signaling , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Production of Molecular Mediator of Immune Response , Proton Transport , Activated T Cell Proliferation

Application Details

Application Notes:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL, 15.6pg/mL human IL-4
--------------------	--

Application Details

standard solutions into the pre-coated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each human IL-4 standard solution and each sample be measured in duplicate.

Comment: Standard: Expression system for standard: E.coli, Immunogen sequence: H25-S153

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only


Handling

Storage: 4 °C,-20 °C

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiry Date: 12 months

Images



ELISA

Image 1. Human IL-4 Accusignal ELISA Kit Human IL-4 Accusignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <1.5pg/ml.