antibodies .- online.com







Osteoactivin ELISA Kit



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	1 kit
Target:	Osteoactivin (GPNMB)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	156 pg/mL - 10000 pg/mL
Minimum Detection Limit:	156 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen	
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Synonyms: Gpnmb, GPNMB_HUMAN, Transmembrane glycoprotein HGFIN, Transmembrane glycoprotein NMB	

Background: Transmembrane glycoprotein NMB is a protein that is encoded by the GPNMB gene. In osteoblast progenitor cells, GPNMB works as a positive regulator of osteoblast differentiation during later stages of matrix maturation and mineralization that is mediated at least in part by BMP-2 in a SMAD1 dependent manner to promote osteoblast differentiation. GPNMB can enhance the repairing process in bone fracture, demonstrated by its high expression during chondrogenesis(soft callus) and osteogenesis(hard callus) compared to the intact femurs that is why Osteoactivin(OA) could be a novel therapeutic agent used to treat generalized osteoporosis or localized osteopenia during fracture repair by stimulating bone growth and regeneration. Similarly, GPNMB expression increases during osteoclast differentiation and it is functionally implicated in this process, possibly by promoting the fusion of osteoclast progenitor cells.

Gene Name: GPNMB

Production: Natural and recombinant mouse Osteoactivin. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: K23-N502

Target Details

Target:	Osteoactivin (GPNMB)	
Alternative Name:	Osteoactivin - GPNMB (GPNMB Products)	
Gene ID:	93695	
NCBI Accession:	NP_444340	
UniProt:	Q99P91	

Application Details

Application	Notes:
-------------	--------

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 10,000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 313pg/mL, 156pg/mL mouse Osteoactivin standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. We recommend that each mouse Osteoactivin standard solution and each sample is measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 156pg/mL-10,000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only

Handling

Storage:	RT,4 °C,-20 °C	
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at	
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only	
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid	
	cycles of freezing and thawing.	