

# Datasheet for ABIN6720099

## CD36 (SR-B3) ELISA Kit





#### Overview

Quantity:	1 kit
Target:	CD36 (SR-B3)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	156 pg/mL - 10000 pg/mL
Minimum Detection Limit:	156 pg/mL
Application:	ELISA

#### **Product Details**

Purpose:	Human CD36 - SR-B3 Sandwich ELISA Kit for Quantitative Detection
Brand:	AccuSignal™
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Production: Natural and recombinant human CD36. There is no detectable cross-reactivity with other relevant proteins.
Sensitivity:	< 10 pg/mL
Components:	<ul><li>Antibody-coated 96-well plate</li><li>Target Protein Standard</li></ul>

Detection antibody

- · Detection reagent
- · Diluent buffers
- · Wash buffers
- · Substrate Solution
- Stop solutions
- · Adhesive covers

#### **Target Details**

Alternative Name:	CD36/SR-B3
Background:	Synonyms: CD36 antigen, CD36 antigen (collagen type I receptor thrombospondin receptor),
	CD36 Molecule, CD36 Molecule (thrombospondin receptor), CHDS7, Cluster determinant 36,
	FAT, Fatty acid translocase, Glycoprotein IIIb, GP3B, GP4, GPIIIB, GPIV, Leukocyte differentiation
	antigen CD36, PAS 4 protein, PAS IV, PASIV, Platelet collagen receptor, Platelet glycoprotein 4,
	Platelet glycoprotein IV, SCARB3, Scavenger receptor class B member 3, Thrombospondin
	receptor
	Background: CD36(Cluster of Differentiation 36) is an integral membrane protein found on the
	surface of many cell types in vertebrate animals and is also known as FAT, SCARB3, GP88,
	glycoprotein IV(gpIV) and glycoprotein IIIb(gpIIIb). The human CD36 is a member of a gene
	family of structurally related glycoproteins and functions as a receptor for collagen type I and
	thrombospondin. The use of a CD36 genomic probe has allowed the localization of the CD36
	locus to the 7q11.2 band by fluorescence in situ hybridization coupled with GTG-banding. CED
	1/SCARF1 and C03F11.3/CD36 are beta-glucan binding receptors and define an evolutionarily
	conserved pathway for the innate sensing of fungal pathogens.
Gene ID:	948
NCBI Accession:	NP_000063
UniProt:	P16671

#### **Application Details**


#### Application Notes:

Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 10000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 312pg/mL, 156pg/mL human CD36 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended

## **Application Details**

	that each human CD36 standard solution and each sample be measured in duplicate.
Comment:	Standard: Expression system for standard: sf21, Immunogen sequence: G30-N439
Sample Volume:	100 μL
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiry Date:	12 months

### **Images**



#### **ELISA**

**Image 1.** Human CD36/SR-B3 Accusignal ELISA Kit Human CD36/SR-B3 AccuSignal ELISA Kit standard curve. Assay Range: 156pg/ml-10000pg/ml. Sensitivity: <10pg/ml.