# antibodies - online.com





## Datasheet for ABIN6720126

### **FCER2 ELISA Kit**



#### Overview

Quantity:	1 kit
Target:	FCER2
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	156 pg/mL - 10000 pg/mL
Minimum Detection Limit:	156 pg/mL
Application:	ELISA

#### Product Details

Product Details	
Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Synonyms: BLAST 2, BLAST-2, C-type lectin domain family 4 member J, CD23, CD23 antigen,
	CD23A, CLEC4J, Fc fragment of IgE low affinity II receptor for (CD23), Fc-epsilon-RII, FCE2,
	Fcer2, FCER2_HUMAN, IGEBF, Immunoglobulin E-binding factor, Low affinity immunoglobulin
	epsilon Fc receptor, Low affinity immunoglobulin epsilon Fc receptor soluble form, Lymphocyte
	IgE receptor
	Background: CD23, also known as Fc epsilon RII, or FceRII, is the "low-affinity" receptor for IgE,
	an antibody isotype involved in allergy and resistance to parasites, and is important in
	regulation of IgE levels. There are two forms of CD23: CD23a and CD23b. CD23a is present on

follicular B cells, whereas CD23b requires IL-4 to be expressed on T-cells, monocytes, Langerhans cells, eosinophils, and macrophages. As part of a mapping of multiple probes to specific bands on chromosome 19 by fluorescence in situ hybridization, the CD23 gene was assigned to 19p13.3. And CD23 (FCE2) is a key molecule for B-cell activation and growth. It is the low-affinity receptor for IgE. The truncated molecule can be secreted, then functioning as a potent mitogenic growth factor.

Gene Name: CD23

Production: Natural and recombinant mouse CD23. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: E50-P331

#### **Target Details**

Target:	FCER2
Alternative Name:	CD23 - FCER2 (FCER2 Products)
Gene ID:	12759
NCBI Accession:	NP_038520
UniProt:	Q06890
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process

#### **Application Details**

<b>Application Notes</b>	daA	lication	Notes:
--------------------------	-----	----------	--------

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 10000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 312pg/mL, 156pg/mL mouse CD23 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each mouse CD23 standard solution and each sample be measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 156pg/mL-10000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only

## Handling

Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid
	cycles of freezing and thawing.