# antibodies -online.com





# **CD137 ELISA Kit**





#### Overview

Quantity:	1 kit
Target:	CD137 (TNFRSF9)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

#### **Product Details**

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Synonyms: 4-1BB, 4-1BB ligand receptor, CD137, CD137 antigen, CDw137, Homolog of mouse
	4 1BB, ILA, Induced by lymphocyte activation (ILA), Interleukin activated receptor homolog of
	mouse Ly63, Receptor protein 4 1BB, T-cell antigen 4-1BB homolog, T-cell antigen ILA,
	TNFRSF9, TNR9_HUMAN, Tumor necrosis factor receptor superfamily member 9
	Background: CD137 is a member of the tumor necrosis factor(TNF) receptor family. Its
	alternative names are tumor necrosis factor receptor superfamily member 9(TNFRSF9), 4-1BB
	and induced by lymphocyte activation(ILA). CD137 gene was localized to chromosome
	1p36.Members of this receptor family and their structurally related ligands are important

regulators of a wide variety of physiological processes and play an especially important role in the regulation of immune responses. CD137 can be expressed by activated T cells, but to a larger extent on CD8 than on CD4 T cells. The best characterized activity of CD137 is its costimulatory activity for activated T cells. Crosslinking of CD137 enhances T cell proliferation, IL-2 secretion survival and cytolytic activity. Further, it can enhance immune activity to eliminate tumors in mice.

Gene Name: TNFRSF9

Production: Natural and recombinant human TNFRSF9. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: L24-Q186

### **Target Details**

Target:	CD137 (TNFRSF9)
Alternative Name:	TNFRSF9 - 4-1BB (TNFRSF9 Products)
Gene ID:	3604
NCBI Accession:	NP_001552
UniProt:	Q07011
Pathways:	Cancer Immune Checkpoints

## **Application Details**

Application Notes:	Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot
	0.1 mL per well of the 2000pg/mL,1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL,

31.2pg/mL human TNFRSF9 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well.

It is recommended that each human TNFRSF9 standard solution and each sample be

measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 31.2pg/mL-2000pg/mL

Sample Volume: 100 µL


Plate: Pre-coated

Restrictions: For Research Use only

### Handling

Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid
	cycles of freezing and thawing.

# **Images**



#### **ELISA**

Image 1. Human TNFRSF9/4-1BB Accusignal ELISA Kit Human TNFRSF9/4-1BB AccuSignal ELISA Kit standard curve. Assay Range: 31.2pg/ml-2000pg/ml. Sensitivity: <10pg/ml.