antibodies .- online.com





F11R ELISA Kit

Overview

Quantity:	1 kit
Target:	F11R
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	31.2 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.2 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen	
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Synonyms: CD 321, CD321, CD321 antigen, ESTM33, F11 receptor, F11R, JAM 1, JAM A, JAM, JAM-1, JAM-A, JAM1, JAM1, JAM1, JAM1, JAM1, JCAM, JCAM, Jcam1, Junction adhesion molecule 1, Junction adhesion molecule, mouse, homolog of, Junctional adhesion molecule 1, Junctional adhesion molecule A, KAT, Ly106, PAM 1, PAM-1, PAM1, Platelet adhesion molecule 1, Platelet adhesion molecule, Platelet F11 receptor, PRO301, UNQ264, UNQ264/PRO301	

Background: Junctional adhesion molecule A(JAM-A) is a protein that in humans is encoded by the F11R gene. It is mapped to 1q23.3. This gene is an immunoglobulin-like molecule that colocalizes with tight junctions in endothelium and epithelium and is also found on blood

leukocytes and platelets. JAM-A plays an important role in the regulation of tight junction assembly in epithelia. In addition, it can act as a receptor for reovirus, a ligand for the integrin LFA1, involved in leukocyte transmigration, and a platelet receptor. JAM-A has a nonredundant role in controlling DC motility, trafficking to lymph nodes, and activation of specific immunity. Gene Name: F11R

Production: Natural and recombinant mouse JAM-A. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: K27-G238

Target Details

Target:	F11R
Alternative Name:	JAM-A - F11R (F11R Products)
Gene ID:	16456
NCBI Accession:	NP_766235
UniProt:	088792
Pathways:	Cell-Cell Junction Organization

Application Details

Application Notes:	Applica	ation Notes:	Α
--------------------	---------	--------------	---

pplication Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL mouse JAM-A standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each mouse JAM-A standard solution and each sample be measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 31.2pg/mL-2000pg/mL

Sample Volume: 100 μL

Plate: Pre-coated

Restrictions: For Research Use only

Handling

Storage:	RT,4 °C,-20 °C	
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at	
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only	
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid	
	cycles of freezing and thawing.	