



[Go to Product page](#)

Datasheet for ABIN6720234
TGFBI ELISA Kit

Overview

Quantity:	1 kit
Target:	TGFBI
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	62.5 pg/mL - 4000 pg/mL
Minimum Detection Limit:	62.5 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	<p>Synonyms: Beta ig-h3, beta-IG-H3, BIGH3, CDB1, CDG2, CDGG1, CSD, CSD1, CSD2, CSD3, EBMD, Kerato-epithelin, LCD1, RGD-CAP, RGD-containing collagen-associated protein, TGFBI, Transforming growth factor-beta-induced protein ig-h3</p> <p>Background: Transforming growth factor, beta-induced, 68 kDa, also known as TGFBI(initially called BIGH3, BIG-H3), is a protein which in humans is encoded by the TGFBI gene. It is mapped to 5q31.1. This gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-</p>

Product Details

collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion.

Gene Name: TGFBI

Production: Natural and recombinant human TGFBI. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: G24-H683

Target Details

Target: TGFBI

Alternative Name: TGFBI - beta IG-H3 ([TGFBI Products](#))

Gene ID: 7045

NCBI Accession: [NP_000349](#)

UniProt: [Q15582](#)

Application Details

Application Notes: Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 4000pg/mL, 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL human TGFBI standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each human TGFBI standard solution and each sample be measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 62.5pg/mL-4000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only

Handling

Storage: RT, 4 °C, -20 °C

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only

Handling

prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.