

Datasheet for ABIN6720255
TNFRSF18 ELISA Kit[Go to Product page](#)

1 Image

Overview

Quantity:	1 kit
Target:	TNFRSF18
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	62.5 pg/mL - 4000 pg/mL
Minimum Detection Limit:	62.5 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric

Characteristics:	<p>Synonyms: Activation inducible TNFR family receptor, AITR, GITR, GITR D, Glucocorticoid induced TNFR related protein, Tumor necrosis factor receptor superfamily member 19</p> <p>Background: Tumor necrosis factor receptor superfamily member 18(TNFRSF18), also called GITR or AITR is a protein that in humans is encoded by the TNFRSF18 gene. This gene is mapped to 1p36.33. This gene encodes a member of the TNF-receptor superfamily. The encoded receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. Knockout studies in mice also suggest the role of this</p>
------------------	--

Product Details

receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Gene Name: TNFRSF18

Production: Natural and recombinant mouse TNFRSF18. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: S22-H153

Target Details

Target:	TNFRSF18
Alternative Name:	TNFRSF18 - GITR (TNFRSF18 Products)
Gene ID:	21936
NCBI Accession:	NP_033426
UniProt:	O35714
Pathways:	Cancer Immune Checkpoints

Application Details

Application Notes:	<p>Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 4000pg/mL, 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL mouse TNFRSF18 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each mouse TNFRSF18 standard solution and each sample be measured in duplicate.</p> <p>Blood Product Anticoagulant: Heparin Sodium</p> <p>ELISA Dilution: 62.5pg/mL-4000pg/mL</p>
Sample Volume:	100 µL
Plate:	Pre-coated
Restrictions:	For Research Use only

Handling

Storage:	RT, 4 °C, -20 °C
----------	------------------

Handling

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images

