

[Go to Product page](#)

Datasheet for ABIN6720269

MME ELISA Kit

Overview

Quantity:	1 kit
Target:	MME
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	93.7 pg/mL - 6000 pg/mL
Minimum Detection Limit:	93.7 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	<p>Synonyms: Atriopeptidase, CALLA, CD10, CD10 antigen, Common acute lymphocytic leukemia antigen, Enkephalinase, EPN, gp100, Membrane metallo endopeptidase, Membrane metallo endopeptidase variant 1, Membrane metallo endopeptidase variant 2, Membrane metalloendopeptidase, Membrane metalloendopeptidase neutral endopeptidase enkephalinase, Membrane metalloendopeptidase variant 1, Membrane metalloendopeptidase variant 2, NEP, NEP_HUMAN, Neprilysin, Neutral endopeptidase, SFE, Skin fibroblast elastase</p> <p>Background: Neprilysin, also known as membrane metallo-endopeptidase, neutral endopeptidase(NEP), CD10, and common acute lymphoblastic leukemia antigen(CALLA), is a</p>

Product Details

zinc-dependent metalloprotease enzyme that degrades a number of small secreted peptides, most notably the amyloid beta peptide whose abnormal misfolding and aggregation in neural tissue has been implicated as a cause of Alzheimer's disease, this gene is localized to human chromosome 3 by study of somatic cell hybrids and regionalized the location to 3q21-q27 by in situ hybridization. By cDNA transfection analysis, Neprilysin is confirmed as a functional neutral endopeptidase of the type that has previously been called enkephalinase. Neprilysin has also been called atriopeptidase. Atriopeptidase specifically degrades atrial natriuretic factor. A specific enzyme inhibitor was developed and reported that it had effects similar to those of low-dose ANF infusion. These effects include diuresis, natriuresis, vasodilatation, and suppression of the renin-angiotensin-aldosterone system.

Gene Name: MME

Production: Natural and recombinant mouse CD10. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: Y52-W750

Target Details

Target:	MME
Alternative Name:	Neprilysin - CD10 (MME Products)
Target Type:	Chemical
Gene ID:	17380
NCBI Accession:	NP_001276391
UniProt:	Q61391
Pathways:	RTK Signaling , Peptide Hormone Metabolism , Regulation of Systemic Arterial Blood Pressure by Hormones , Smooth Muscle Cell Migration

Application Details

Application Notes:	Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 6000pg/mL, 3000pg/mL, 1500pg/mL, 750pg/mL, 419pg/mL, 187.5pg/mL, 93.7pg/mL mouse CD10 standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each mouse CD10 standard solution and each sample be measured in duplicate.
--------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Application Details

Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 93.7pg/mL-6000pg/mL

Sample Volume: 100 µL

Plate: Pre-coated

Restrictions: For Research Use only

Handling

Storage: RT, 4 °C, -20 °C

Storage Comment: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.