# antibodies .- online.com







#### **PD-L1 ELISA Kit**



#### Overview

Quantity:	1 kit
Target:	PD-L1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	62.5 pg/mL - 4000 pg/mL
Minimum Detection Limit:	62.5 pg/mL
Application:	ELISA

#### **Product Details**

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Synonyms: B7 H, B7H, B7 H1, B7-H1, B7H1, CD 274, CD274, PD L1, PD-L1, PDL 1, PDL1  Background: Programmed death-ligand 1 (PD-L1), also known as cluster of differentiation 274

Background: Programmed death-ligand 1 (PD-L1), also known as cluster of differentiation 274 (CD274) or B7 homolog 1 (B7-H1), is a protein that in humans is encoded by the CD274 gene. PD-L1 is a 40 kDa type 1 transmembrane protein that has been speculated to play a major role in suppressing the immune system during particular events such as pregnancy, tissue allografts, autoimmune disease and other disease states such as hepatitis. Normally the immune system reacts to foreign antigens where there is some accumulation in the lymph nodes or spleen which triggers a proliferation of antigen-specific CD8+ T cell. The formation of PD-1 receptor / PD-L1 or B7.1 receptor /PD-L1 ligand complex transmits an inhibitory signal which reduces the proliferation of these CD8+ T cells at the lymph nodes and supplementary to that PD-1 is also able to control the accumulation of foreign antigen specific T cells in the lymph nodes through apoptosis which is further mediated by a lower regulation of the gene Bcl-2.

Gene Name: PD-L1

Production: Natural and recombinant human PD-L1. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: F19-R238

### **Target Details**

Target:	PD-L1
Alternative Name:	PD-L1 - B7-H1 (PD-L1 Products)
Gene ID:	2848
NCBI Accession:	NP_001254635
UniProt:	Q9NZQ7
Pathways:	Cancer Immune Checkpoints

Application Details	
Application Notes:	Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot
	0.1 mL per well of the 4000pg/mL, 2000pg/mL,1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL,
	62.5pg/mL human PD-L1 standard solutions into the precoated 96-well plate. Add 0.1 mL of the
	sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted
	sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well.
	It is recommended that each human PD-L1 standard solution and each sample be measured in
	duplicate.
	Blood Product Anticoagulant: Heparin Sodium
	ELISA Dilution: 62.5pg/mL-4000pg/mL
Sample Volume:	100 μL
Plate:	Pre-coated
Restrictions:	For Research Use only

## Handling

Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at
	room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only
	prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid
	cycles of freezing and thawing.