

Datasheet for ABIN6720311

CLEC3B ELISA Kit



Overview

Quantity:	1 kit
Target:	CLEC3B
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	312 pg/mL - 20000 pg/mL
Minimum Detection Limit:	312 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen	
Sample Type:	Cell Culture Supernatant, Serum, Tissue Homogenate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Synonyms: C type lectin domain family 3 member B, C-type lectin domain family 3 member B,	

Synonyms: C type lectin domain family 3 member B, C-type lectin domain family 3 member B, Clec3b, Plasminogen kringle 4 binding protein, Plasminogen kringle 4-binding protein, TETN_HUMAN, Tetranectin (plasminogen binding protein), Tetranectin, TN, TNA Background: Tetranectin, also called TNA, is a protein that in humans is encoded by the CLEC3B gene. It is mapped to 3p21.31. Tetranectin, a tetrameric protein isolated from human plasma, has 4 identical and noncovalently bound polypeptide chains, each of 181 amino acid residues. It has a specific binding affinity for sulfated polysaccharides and the kringle 4 of plasminogen. The plasma concentration of tetranectin is reduced in patients with various

malignancies. Tetranectin is a plasminogen-binding protein that is induced during the mineralization phase of osteogenesis. Thus, tetranectin is a candidate gene for human disorders affecting bone and connective tissue.

Gene Name: CLEC3B

Production: Natural and recombinant human Tetranectin . There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: NSO, Immunogen sequence: E22-V202

Target Details

Target:	CLEC3B	
Alternative Name:	Tetranectin - CLEC3B (CLEC3B Products)	
Gene ID:	7123	
NCBI Accession:	NP_003269	
UniProt:	P05452	

Application Details

۸۰۰	lication	Niataa
Ann	lication	INOTES:

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 20,000pg/mL, 10,000pg/mL, 5000pg/mL, 2500pg/mL, 1250pg/mL, 625pg/mL, 312pg/mL human Tetranectin standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of human cell culture supernates, serum or tissue Homogenates to each empty well. We recommend that each human Tetranectin standard solution and each sample is measured in duplicate.

ELISA Dilution: 312pg/mL-20,000pg/mL

Sample Volume:

100 μL

Plate:

Pre-coated

Restrictions:

For Research Use only

Handling

O+	RT.4 °C20 °C
Storage:	R I 4 U/II U.

Storage Comment:

Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only

prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.