antibodies -online.com







ACVA ELISA Kit





Overview

Quantity:	1 kit
Target:	ACVA
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	15.6 pg/mL - 1000 pg/mL
Minimum Detection Limit:	15.6 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Saliva, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	Synonyms: Activin beta A chain, erythroid differentiation factor, erythroid differentiation protein, follicle stimulating hormone releasing protein, FSH releasing Background: Activin A is a homodimer of 14 kDa beta-A. Activin A, a cytokine member of the

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn

transforming growth factor-beta superfamily, is expressed locally by the mesenchymal

component of the hemopoietic microenvironment. Its expression is regulated on the mRNA

level by different cytokines, and the biological activity of the protein is tightly controlled by

several inhibitory molecules. Inhibins and activins are members of the transforming growth factor beta superfamily and are known to modulate the growth and differentiation of several cell types. Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins appear to oppose the functions of activins. The standard product used in this kit is recombinant Activin A, which is composed of two single chains of 116 amino acids with the molecular mass of 26KDa.

Gene Name: INHBA

Production: Natural and recombinant mouse Activin A. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: CHO, Immunogen sequence: G311-S427

Target Details

Target:	ACVA
Alternative Name:	Activin A (ACVA Products)
Gene ID:	16323
NCBI Accession:	NP_032406
UniProt:	Q04998
Pathways:	Hormone Transport, Peptide Hormone Metabolism

Application Details

Application Notes:

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL, 15.6pg/mL mouse Activin A standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of mouse cell culture supernates, serum, plasma(heparin, EDTA) or saliva to each empty well. It is recommended that each mouse Activin A standard solution and each sample be measured in duplicate.

Blood Product Anticoagulant: Heparin Sodium


ELISA Dilution: 15.6pg/mL-1000pg/mL

Sample Volume: 100 µL

Application Details

Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Images



ELISA

Image 1. Mouse Activin A Accusignal ELISA KIT Mouse Activin A AccuSignal Elisa Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Senstivity: <10pg/ml.