# antibodies .- online.com







## **ACVA ELISA Kit**





## Overview

Quantity:	1 kit
Target:	ACVA
Reactivity:	Rat
Method Type:	Sandwich ELISA
Detection Range:	15.6 pg/mL - 1000 pg/mL
Minimum Detection Limit:	15.6 pg/mL
Application:	ELISA

# **Product Details**

Purpose:	Sandwich ELISA for Quantitative Detection of Antigen	
Sample Type:	Cell Culture Supernatant, Plasma (EDTA - heparin), Saliva, Serum	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Synonyms: Activin beta A chain, erythroid differentiation factor, erythroid differentiation protein, follicle stimulating hormone releasing protein, FSH releasing	
	Background: Activin A is a homodimer of 14 kDa beta-A. Activin A, a cytokine member of the	

transforming growth factor-beta superfamily, is expressed locally by the mesenchymal component of the hemopoietic microenvironment. Its expression is regulated on the mRNA level by different cytokines, and the biological activity of the protein is tightly controlled by several inhibitory molecules. Inhibins and activins are members of the transforming growth factor beta superfamily and are known to modulate the growth and differentiation of several cell types. Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins appear to oppose the functions of activins. The standard product used in this kit is recombinant Activin A, which is composed of two single chains of 116 amino acids with the molecular mass of 26KDa.

Gene Name: INHBA

Production: Natural and recombinant rat Activin A. There is no detectable cross-reactivity with other relevant proteins.

Standard: Expression system for standard: CHO, Immunogen sequence: G311-S428

## **Target Details**

Target:	ACVA
Alternative Name:	Activin A (ACVA Products)
Gene ID:	29200
NCBI Accession:	NP_058824
UniProt:	P18331
Pathways:	Hormone Transport, Peptide Hormone Metabolism

## **Application Details**

#### **Application Notes:**

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1 mL per well of the 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL, 15.6pg/mL rat Activin A standard solutions into the precoated 96-well plate. Add 0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each properly diluted sample of rat cell culture supernates, serum, plasma(heparin, EDTA) or saliva to each empty well. It is recommended that each rat Activin A standard solution and each sample be measured in duplicate. Blood Product Anticoagulant: Heparin Sodium

ELISA Dilution: 15.6pg/mL-1000pg/mL

100 μL

Sample Volume:

## **Application Details**

Plate:	Pre-coated
Restrictions:	For Research Use only


# Handling

Storage:	RT,4 °C,-20 °C	

Storage Comment:

Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

## **Images**



### **ELISA**

**Image 1.** Rat Activin A Accusignal ELISA Kit Rat Activin A AccuSignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <12pg/ml.