

Datasheet for ABIN6723489

anti-F4/80 antibody (Alexa Fluor 647)



Go to Product page

| _ | | | | | |
|---|---|---|----|----|---|
| | W | 0 | rv | 10 | W |

| Quantity: | 100 μg | |
|-------------------|--|--|
| Target: | F4/80 (EMR1) | |
| Reactivity: | Mouse | |
| Host: | Rat | |
| Clonality: | Monoclonal | |
| Conjugate: | This F4/80 antibody is conjugated to Alexa Fluor 647 | |
| Application: | Flow Cytometry (FACS) | |
| | | |
| Product Details | | |
| Clone: | CI:A3-1 | |
| Isotype: | IgG2b kappa | |
| Target Details | | |
| Target: | F4/80 (EMR1) | |
| Alternative Name: | F4/80 (EMR1 Products) | |
| Background: | Synonyms: Adhesion G protein-coupled receptor E1,Adgre1,Cell surface glycoprotein | |
| | F4/80,EGF-like module receptor 1,Adgre1,Emr1, Gpf480 | |
| | Background: F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal | |
| | growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has | |
| | been widely used as a murine macrophage marker, which is expressed on the majority of tissue | |
| | macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red | |

Target Details

| pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node |
|--|
| and Peyer's patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has |
| also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been |
| identified, but it has been reported that F4/80 is required for induction of CD8+ T cells-mediated |
| peripheral tolerance. |
| |

Gene ID:

13733

UniProt:

Q61549

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. | |
|--------------------|--|--|
| Restrictions: | For Research Use only | |

Handling

| Concentration: | 0.5 mg/mL | |
|--------------------|--|--|
| Buffer: | PBS with 0.05 % Proclin300, 1 % BSA | |
| Preservative: | ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Precaution of Use: | | |
| Storage: | 4 °C | |
| Storage Comment: | Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. | |