# antibodies -online.com





## Datasheet for ABIN6730244

## Olfactomedin 1 ELISA Kit



#### Overview

Quantity:	96 tests
Target:	Olfactomedin 1 (OLFM1)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	0.084 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.084 ng/mL
Application:	ELISA

#### **Product Details**

Purpose:	Mouse Olfactomedin-1 ELISA Kit for Serum, Plasma, and Cell Culture Supernatants.
Sample Type:	Cell Culture Supernatant, Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair recognizes Mouse Olfactomedin-1.
Characteristics:	Strip plates and additional reagents allow for use in multiple experiments
	Quantitative protein detection
	Establishes normal range
	The best products for confirmation of antibody array data
Components:	Pre-Coated 96-well Strip Microplate
	Wash Buffer
	Stop Solution

#### **Product Details**

- Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- · Tubes to prepare standard and sample dilutions
- · Absorbent paper

Olfactomedin 1 (OLFM1)

- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

### **Target Details**

Target:

Alternative Name:	Olfactomedin-1 (OLFM1 Products)	
UniProt:	O88998	
Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.	
	2. Add 100 µl of standard or sample to each well.	
	3. Incubate 2.5 h at RT or O/N at 4°C.	
	4. Add 100 μl of prepared biotin antibody to each well.	
	5. Incubate 1 h at RT.	
	6. Add 100 μl of prepared Streptavidin solution to each well.	
	7. Incubate 45 min at RT.	
	8. Add 100 µl of TMB One-Step Substrate Reagent to each well.	
	9. Incubate 30 min at RT.	
	10. Add 50 μl of Stop Solution to each well.	
	11. Read at 450 nm immediately.	
Reagent Preparation:	Recommended Dilution for serum and plasma samples10 fold	
Restrictions:	For Research Use only	

## Handling

Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated
	freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is
	recommended to store at -80°C.
Expiry Date:	6 months