# antibodies -online.com





# Datasheet for ABIN6730296

# **SDC4 ELISA Kit**



# Overview

Quantity:	96 tests
Target:	SDC4
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

### Product Details

Product Details					
Purpose:	Mouse SDC4 ELISA Kit.				
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate				
Analytical Method:	Quantitative				
Detection Method:	Colorimetric				
Specificity:	This ELISA antibody pair recognizes Mouse SDC4.				
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>				
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> </ul>				

# **Product Details**

- · Streptavidin-Conjugated HRP
- · TMB One-Step Substrate

### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- · Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

# **Target Details**

Target:	SDC4
Alternative Name:	SDC4 (SDC4 Products)
UniProt:	035988
Pathways:	Glycosaminoglycan Metabolic Process

# **Application Details**

10 10 10 10 10 10 10 10 10 10 10 10 10 1				
Application Notes:	Optimal working dilution should be determined by the investigator.			
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.			
	2. Add 100 µl of standard or sample to each well.			
	3. Incubate 2.5 h at RT or O/N at 4°C.			
	4. Add 100 µl of prepared biotin antibody to each well.			
	5. Incubate 1 h at RT.			
	6. Add 100 µl of prepared Streptavidin solution to each well.			
	7. Incubate 45 min at RT.			
	8. Add 100 µl of TMB One-Step Substrate Reagent to each well.			
	9. Incubate 30 min at RT.			
	10. Add 50 µl of Stop Solution to each well.			
	11. Read at 450 nm immediately.			
Restrictions:	For Research Use only			
Handling				
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated			
	freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is			

1	1		11.		
_	$\sim$	$\sim$	4 I I	n	$\sim$
_	14	111	111	1 1	

recommended to store at -80°C.

Expiry Date: 6 months