# antibodies -online.com







## **CTHRC1 ELISA Kit**



( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	96 tests
Target:	CTHRC1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	8 pg/mL - 5000 pg/mL
Minimum Detection Limit:	8 pg/mL
Application:	ELISA

Product Details	
Purpose:	Human CTHRC1 ELISA Kit for Serum, Plasma, and Cell Culture Supernatants.
Sample Type:	Cell Culture Supernatant, Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair recognizes Human CTHRC1.
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>
Components:	<ul><li>Pre-Coated 96-well Strip Microplate</li><li>Wash Buffer</li><li>Stop Solution</li></ul>

## **Product Details**

- Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- · TMB One-Step Substrate

#### Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- · Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

## **Target Details**

Target:	CTHRC1
Alternative Name:	CTHRC1 (CTHRC1 Products)
UniProt:	Q96CG8
Pathways:	Tube Formation

## **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.
Protocol:	Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 µl of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4°C.
	4. Add 100 μl of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 µl of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 µl of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μl of Stop Solution to each well.
	11. Read at 450 nm immediately.
Reagent Preparation:	Recommended Dilution for serum and plasma samples5 fold
Restrictions:	For Research Use only

# Handling

Expiry Date:	6 months	
	recommended to store at -80°C.	
	freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is	
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated	