



[Go to Product page](#)

## Datasheet for ABIN6730487

### GAPDH ELISA Kit

#### Overview

Quantity:	96 tests
Target:	GAPDH
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

#### Product Details

Purpose:	Mouse GAPDH ELISA Kit.
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair recognizes Mouse GAPDH.
Characteristics:	<ul style="list-style-type: none"> <li>• Strip plates and additional reagents allow for use in multiple experiments</li> <li>• Quantitative protein detection</li> <li>• Establishes normal range</li> <li>• The best products for confirmation of antibody array data</li> </ul>
Components:	<ul style="list-style-type: none"> <li>• Pre-Coated 96-well Strip Microplate</li> <li>• Wash Buffer</li> <li>• Stop Solution</li> <li>• Assay Diluent(s)</li> <li>• Lyophilized Standard</li> <li>• Biotinylated Detection Antibody</li> </ul>

## Product Details

---

- Streptavidin-Conjugated HRP
- TMB One-Step Substrate

### Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 µl to 1 µl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

## Target Details

---

Target: GAPDH

Alternative Name: GAPDH ([GAPDH Products](#))

UniProt: [P16858](#)

## Application Details

---

Application Notes: Optimal working dilution should be determined by the investigator.

Protocol:

1. Prepare all reagents, samples and standards as instructed in the manual.
2. Add 100 µl of standard or sample to each well.
3. Incubate 2.5 h at RT or O/N at 4°C.
4. Add 100 µl of prepared biotin antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µl of prepared Streptavidin solution to each well.
7. Incubate 45 min at RT.
8. Add 100 µl of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µl of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only

## Handling

---

Storage Comment: The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.

## Handling

---

Expiry Date: 6 months