

Datasheet for ABIN6730568

ABL1 ELISA Kit**3** Images[Go to Product page](#)

Overview

Quantity: 96 tests

Target: ABL1

Binding Specificity: pTyr245

Reactivity: Human

Method Type: Sandwich ELISA

Application: ELISA

Product Details

Purpose: Human Phospho-c-Abl (Tyr245) and Total c-Abl ELISA Kit. This assay semi-quantitatively measures c-Abl phosphorylated at Tyrosine-245 as well as total c-Abl in cell lysate samples.

Sample Type: Cell Samples, Tissue Lysate

Analytical Method: Semi-Quantitative

Detection Method: Colorimetric

Specificity: This ELISA kit recognizes Human c-Abl phosphorylated at site Tyrosine-245 as well as total c-Abl.

Characteristics:

- Pre-Coated 96-well Strip Microplate
- Wash Buffer
- Anti-Phospho Antibody
- Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- Streptavidin-Conjugated HRP
- Assay Diluent

Product Details

- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Components:

- Pre-Coated 96-well Strip Microplate
- Wash Buffer
- Anti-Phospho Antibody
- Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- Streptavidin-Conjugated HRP
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: ABL1

Alternative Name: c-Abl ([ABL1 Products](#))

Gene ID: 25

UniProt: [P00519](#)

Pathways: [Apoptosis](#), [Regulation of Muscle Cell Differentiation](#), [Platelet-derived growth Factor Receptor Signaling](#), [Lipid Metabolism](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Protocol: 1. Prepare all reagents and samples as instructed in the manual.

Application Details

2. Add 100 μ L of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 $^{\circ}$ C.
4. Add 100 μ L of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 μ L of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: -20 $^{\circ}$ C

Storage Comment: Upon receipt, the kit should be stored at -20 $^{\circ}$ C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 $^{\circ}$ C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 $^{\circ}$ C. Reconstituted Positive Control (Item K) should be stored at -70 $^{\circ}$ C.


Expiry Date: 6 months

Images




ELISA

Image 1. Jurkat cells were treated with Pervanadate. Cell lysates were analyzed using this phosphoELISA and Western Blot.



ELISA

Image 2. Jurkat cells were treated with Pervanadate. Solubilize cells at 4×10^7 cells/ml in Cell Lysate Buffer. Serial dilutions of lysates were analyzed in this ELISA.



ELISA

Image 3. Jurkat cells were treated with Pervanadate. Cell lysates were analyzed using this phosphoELISA and Western Blot.