antibodies .- online.com





Datasheet for ABIN6730593

FLT3 ELISA Kit

3 Images



Go to Product page

Overview

Quantity:	96 tests
Target:	FLT3
Binding Specificity:	pTyr589
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA
Product Details	
Purpose:	Human Phospho-FLT3 (Tyr589) ELISA Kit. This assay semi-quantitatively measures FLT3 phosphorylated at Tyrosine-589 in cell lysate samples.
Sample Type:	Cell Samples, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human FLT3 phosphorylated at site Tyrosine-589.
Characteristics:	 Pre-Coated 96-well Strip Microplate Wash Buffer Anti-Phospho Antibody HRP-Conjugated Secondary Antibody Assay Diluent TMB One-Step Substrate Stop Solution Lysis Buffer

Product Details

	Positive Control Sample	
Components:	Pre-Coated 96-well Strip Microplate	
	Wash Buffer	

- Anti-Phospho Antibody
- · HRP-Conjugated Secondary Antibody
- · Assay Diluent
- TMB One-Step Substrate
- · Stop Solution
- · Lysis Buffer
- · Positive Control Sample

Material not included:

- · Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- · Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	FLT3
Alternative Name:	FLT3 (FLT3 Products)
Gene ID:	2322
UniProt:	P36888
Pathways:	RTK Signaling

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Protocol:	1. Prepare all reagents and samples as instructed in the manual.
	2. Add 100 μ L of sample or positive control to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μ L of prepared primary antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared 1X HRP-Streptavidin to each well.
	7. Incubate 1 h at RT.

- 8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions:

For Research Use only

Handling

Sto	

-20 °C

Storage Comment:


Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date:


6 months

Images

PV




Anti FLT3 (Tyr589)



Anti pan- FLT3


ELISA

Image 1. A431 cells were treated or untreated with Pervanadate. Cell lysates were analyzed using this phosphoELISA and Western Blot.



ELISA

Image 2. Jurkat cells were treated with Pervanadate. Solubilize cells at 4 x 10^7 cells/ml in Cell Lysate Buffer. Serial dilutions of lysates were analyzed in this ELISA.



ELISA

Image 3. A431 cells were treated or untreated with Pervanadate. Cell lysates were analyzed using this phosphoELISA and Western Blot.