antibodies .- online.com





Datasheet for ABIN6730677

FOLR1 ELISA Kit



Overview

Quantity:	96 tests
Target:	FOLR1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details						
Purpose:	Mouse FOLR1 ELISA Kit.					
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate					
Analytical Method:	Quantitative					
Detection Method:	Colorimetric					
Specificity:	This ELISA antibody pair recognizes Mouse FOLR1.					
Characteristics:	 Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data 					
Components:	 Pre-Coated 96-well Strip Microplate Wash Buffer Stop Solution Assay Diluent(s) Lyophilized Standard Biotinylated Detection Antibody 					

Product Details

- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target:	FOLR1
Alternative Name:	FOLR1 (FOLR1 Products)
UniProt:	P35846
Pathways:	Dicarboxylic Acid Transport

Application Details

Application Notes: Optimal working dilution should be determined by the investigator. Protocol: 1. Prepare all reagents, samples and standards as instructed in the manual. 2. Add 100 µl of standard or sample to each well. 3. Incubate 2.5 h at RT or O/N at 4°C. 4. Add 100 µl of prepared biotin antibody to each well. 5. Incubate 1 h at RT.	Application Details				
 Add 100 μl of standard or sample to each well. Incubate 2.5 h at RT or O/N at 4°C. Add 100 μl of prepared biotin antibody to each well. 	Application Notes:				
3. Incubate 2.5 h at RT or O/N at 4°C. 4. Add 100 µl of prepared biotin antibody to each well.	Protocol:				
4. Add 100 μl of prepared biotin antibody to each well.					
5. Incubate 1 h at RT.					
6. Add 100 µl of prepared Streptavidin solution to each well.					
7. Incubate 45 min at RT.					
8. Add 100 μl of TMB One-Step Substrate Reagent to each well.					
9. Incubate 30 min at RT.					
10. Add 50 μl of Stop Solution to each well.					
11. Read at 450 nm immediately.					
Restrictions: For Research Use only	Restrictions:				
Handling	Handling				
Storage Comment: The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repea	Storage Comment:				
freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it					

1	1		11.		
_	\sim	\sim	4 I I	n	\sim
_	14	111	111	1 1	

recommended to store at -80°C.

Expiry Date: 6 months