# antibodies -online.com







## **SPARCL1 ELISA Kit**



( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Quantity:	96 tests
Target:	SPARCL1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details		
Purpose:	Mouse SPARC-like 1/SPARCL1 ELISA Kit.	
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This ELISA antibody pair recognizes Mouse SPARC-like 1.	
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>	
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> </ul>	

#### **Product Details**

- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

### **Target Details**

Target:	SPARCL1	
Alternative Name:	SPARC-like 1 (SPARCL1 (SPARCL1 Products)	
UniProt:	P70663	

Application Details			
Application Notes:	Optimal working dilution should be determined by the investigator.		
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.		
	2. Add 100 µl of standard or sample to each well.		
	3. Incubate 2.5 h at RT or O/N at 4°C.		
	4. Add 100 $\mu$ l of prepared biotin antibody to each well.		
	5. Incubate 1 h at RT.		
	6. Add 100 µl of prepared Streptavidin solution to each well.		
	7. Incubate 45 min at RT.		
	8. Add 100 µl of TMB One-Step Substrate Reagent to each well.		
	9. Incubate 30 min at RT.		
	10. Add 50 µl of Stop Solution to each well.		
	11. Read at 450 nm immediately.		
Restrictions:	For Research Use only		
Handling			
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated		
	freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is		
	recommended to store at -80°C.		

1 1		
$\blacksquare$	$\alpha \alpha \alpha$	lina
	land	1111(
	I G I G	11119

Expiry Date:

6 months