



[Go to Product page](#)

Datasheet for ABIN6731255

IL-2 Protein (AA 21-153) (Fc Tag,AVI tag,Biotin)

3 Images

Overview

Quantity:	200 µg
Target:	IL-2 (IL2)
Protein Characteristics:	AA 21-153
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This IL-2 protein is labelled with Fc Tag,AVI tag,Biotin.

Product Details

Sequence:	AA 21-153
Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	IL-2 (IL2)
Alternative Name:	IL-2 (IL2 Products)
Background:	Interleukin-2 (IL-2) is an interleukin, a type of cytokine immune system signaling molecule,

Target Details

which is a leukocytotropic hormone that is instrumental in the body's natural response to microbial infection and in discriminating between foreign (non-self) and self. IL-2 mediates its effects by binding to IL-2 receptors, which are expressed by lymphocytes, the cells that are responsible for immunity. Mature human IL-2 shares 56 % and 66 % aa sequence identity with mouse and rat IL-2, respectively. Human and mouse IL-2 exhibit crossspecies activity. The receptor for IL-2 consists of three subunits that are present on the cell surface in varying preformed complexes. IL-2 is also necessary during T cell development in the thymus for the maturation of a unique subset of T cells that are termed regulatory T cells (T-regs). After exiting from the thymus, T-Regs function to prevent other T cells from recognizing and reacting against "self antigens", which could result in "autoimmunity". T-Regs do so by preventing the responding cells from producing IL-2. Thus, IL-2 is required to discriminate between self and non-self, another one of the unique characteristics of the immune system.

Molecular Weight: 43.7 kDa

NCBI Accession: [NP_000577](#)

Pathways: [JAK-STAT Signaling](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#), [Activated T Cell Proliferation](#)

Application Details

Comment: Ready-to-use Avitag™ biotinylated protein:
The product is exclusively produced using the Avitag™ technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA.


This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.

Restrictions: For Research Use only

Handling


Format:	Lyophilized
Buffer:	Tris with Glycine, Arginine and NaCl, pH 7.5
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C

Images




ELISA

Image 1. Immobilized Human IL-2 R alpha, Fc Tag (ABIN2181382,ABIN2181381) at 5 µg/mL (100 µL/well) can bind Biotinylated Human IL-2, Fc,Avitag (ABIN6731255,ABIN6809921) with a linear range of 0.6-10 ng/mL (Routinely tested).



ELISA

Image 2. Immobilized Biotinylated Human IL-2, Fc,Avitag (ABIN6731255,ABIN6809921) at 1 µg/mL (100 µL/well) on streptavidin precoated (0.2 µg/well) plate, can bind Human IL-2 R alpha, His Tag (ABIN2181384,ABIN2181383) with a linear range of 1-39 ng/mL (QC tested).



SDS-PAGE

Image 3. Biotinylated Human IL-2, Fc,Avitag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 % .