antibodies .- online.com





anti-SSR1 antibody (AA 51-100)

2 Images

Predicted Reactivity:



Go to i roduct page

Overview	
Quantity:	100 μL
Target:	SSR1
Binding Specificity:	AA 51-100
Reactivity:	Human, Mouse, Dog, Rat, Cow, Horse, Monkey, Rabbit, Zebrafish (Danio rerio), Pig, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SSR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))
Product Details	
Immunogen:	Synthetic peptide located between aa51-100 of human SSR1 (P43307, NP_003135). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan, Gibbon, Monkey, Galago, Marmoset, Mouse, Rat, Elephant, Dog, Bovine, Rabbit, Horse, Pig, Turkey, Zebra finch, Chicken (100%), Guinea pig (92%), Lizard (90%), Xenopus (85%).
	Type of Immunogen: Synthetic peptide
Isotype:	IgG
Specificity:	Human SSR1
Donalista d Danasticita	D

Guinea pig (92%) Xenopus (85%).

Percent identity by BLAST analysis: Human, Mouse, Rat, Dog, Horse, Pig, Chicken (100%)

Product Details Purification: Immunoaffinity purified **Target Details** SSR1 Target: Alternative Name SSR1 (SSR1 Products) Background: Name/Gene ID: SSR1 Synonyms: SSR1, Signal sequence receptor, SSR-alpha, TRAP-alpha, TRAP alpha, Ssr, SSR alpha subunit, Trap-a, TRAPA Gene ID: 6745 NCBI Accession: NP_003135 UniProt: P43307 Pathways: **ER-Nucleus Signaling Application Details** Application Notes: Approved: IHC, IHC-P, WB Usage: ELISA titer using peptide based assay: 1:62500. Western Blot: Suggested dilution at 0.5 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1:50000 - 100000 as second antibody. Target Species of Antibody: Human Comment: Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Distilled Water. Concentration: Lot specific Buffer: Lyophilized from PBS with 2 % sucrose Handling Advice: Avoid repeat freeze-thaw cycles.

4 °C,-20 °C

Storage:

Handling

Storage Comment:

Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long

term use (up to 1 year)

Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.

Images




Image 1.




Image 2.