
antibodies -online.com







anti-TMEM127 antibody (AA 110-159)



Image



Go to	D		
	Pron	ויאווו	mane

_					
U	V	er	V	Ie	W

Quantity:	100 μL
Target:	TMEM127
Binding Specificity:	AA 110-159
Reactivity:	Human, Mouse, Rat, Cow, Rabbit, Dog, Guinea Pig, Pig, Horse, Zebrafish (Danio rerio), Monkey, Bat, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TMEM127 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Synthetic peptide located between aa110-159 of human TMEM127 (075204, NP_060319).
	Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey, Galago,
	Marmoset, Mouse, Rat, Elephant, Panda, Dog, Bovine, Bat, Rabbit, Horse, Pig, Opossum, Guinea
	pig, Xenopus, Salmon, Pike, Pufferfish, Zebrafish (100%).
	Type of Immunogen: Synthetic peptide
Isotype:	IgG
Specificity:	Human TMEM127
Predicted Reactivity:	Percent identity by BLAST analysis: Human, Mouse, Rat, Dog, Bovine, Horse, Xenopus, Zebrafish (100%).

Product Details Purification: Immunoaffinity purified **Target Details** TMEM127 Target: Alternative Name TMEM127 (TMEM127 Products) Background: Name/Gene ID: TMEM127 Synonyms: TMEM127, Transmembrane protein 127 Gene ID: 55654 NCBI Accession: NP 060319 **Application Details** Approved: WB Application Notes: Usage: ELISA titer using peptide based assay: 1:312500. Western Blot: Suggested dilution at 1 µ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1:50000 - 100000 as second antibody. Comment: Target Species of Antibody: Human Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Distilled Water. Concentration: Lot specific Buffer: Lyophilized from PBS with 2 % sucrose Handling Advice: Avoid repeat freeze-thaw cycles. 4 °C,-20 °C Storage: Storage Comment: Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long

Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.

term use (up to 1 year)




Image 1. ransfected 293T cell lysate. Antibody concentration: 1.0 ug/ml. Gel concentration: 12%.