



[Go to Product page](#)

Datasheet for ABIN6740417
anti-KIR5.1 antibody (C-Term)

1 Image

Overview

| | |
|----------------------|---------------------------------------|
| Quantity: | 100 µL |
| Target: | KIR5.1 (KCNJ16) |
| Binding Specificity: | C-Term |
| Reactivity: | Human, Rabbit, Dog, Pig |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This KIR5.1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | Synthetic peptide from C-Terminus of human KCNJ16 (Q9NPI9, NP_061128). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan, Gibbon, Marmoset (100%), Monkey (93%), Panda, Horse (92%), Dog, Rabbit (85%). Type of Immunogen: Synthetic peptide |
| Isotype: | IgG |
| Specificity: | Human KCNJ16 / Kir5.1 |
| Predicted Reactivity: | Percent identity by BLAST analysis: Human (100%) Dog, Rabbit (85%). |
| Purification: | Immunoaffinity purified |

Target Details

| | |
|-------------------|---|
| Target: | KIR5.1 (KCNJ16) |
| Alternative Name: | KCNJ16 / Kir5.1 (KCNJ16 Products) |
| Background: | Name/Gene ID: KCNJ16 Subfamily: Potassium channel - inward-rectifying Family: Ion Channel Synonyms: KCNJ16, BIR9, KIR5.1 |
| Gene ID: | 3773 |
| NCBI Accession: | NP_061128 |
| UniProt: | Q9NPI9 |

Application Details

| | |
|--------------------|--|
| Application Notes: | Approved: WB (0.2 - 1 µg/mL) Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as second antibody. ELISA titer in peptide based assay: 1:12500. |
| Comment: | Target Species of Antibody: Human |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---|
| Format: | Lyophilized |
| Reconstitution: | Distilled water |
| Concentration: | Lot specific |
| Buffer: | Lyophilized from PBS with 2 % sucrose |
| Handling Advice: | Avoid repeat freeze-thaw cycles. |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year) Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles. |




Image 1.