antibodies - online.com







anti-PIM3 antibody (AA 259-288)


Images



\sim					
()	VE	۲۱	/1	\triangle	Λ


Quantity:	200 μL	
Target:	PIM3	
Binding Specificity:	AA 259-288	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PIM3 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Froduct Details		
Brand:	IHC-plus™	
	IHC-plus™ This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic	
Brand:		
Brand:	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic	
Brand: Specificity: Purification:	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 259-288 amino acids from the C-terminal region of human PIM3.	
Brand: Specificity: Purification: Target Details	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 259-288 amino acids from the C-terminal region of human PIM3. Ammonium sulfate precipitation	
Brand: Specificity: Purification: Target Details Target:	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 259-288 amino acids from the C-terminal region of human PIM3. Ammonium sulfate precipitation PIM3	
Brand: Specificity: Purification: Target Details	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 259-288 amino acids from the C-terminal region of human PIM3. Ammonium sulfate precipitation	
Brand: Specificity: Purification: Target Details Target:	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 259-288 amino acids from the C-terminal region of human PIM3. Ammonium sulfate precipitation PIM3	
Brand: Specificity: Purification: Target Details Target: Alternative Name:	This PIM3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 259-288 amino acids from the C-terminal region of human PIM3. Ammonium sulfate precipitation PIM3 PIM3 / PIM-3 (PIM3 Products)	

	Synanyma: DIM2 KID1 Dim 2 anaggana Sarina/thraonina kinaga Dim 2 Dim 2		
	Synonyms: PIM3, KID1, Pim-3 oncogene, Serine/threonine kinase Pim-3, Pim-3		
Gene ID:	415116		
UniProt:	Q86V86		
Pathways:	Negative Regulation of Hormone Secretion		
Application Details			
Application Notes:	Approved: IHC-P (10 μg/mL), WB (1:1000)		
	Usage: Immunohistochemistry: This antibody was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for this antibody was determined to be 10 μ g/mL. Please note that cross-reactivity with mouse has been confirmed in WB only. The antibody has not been evaluated for use in IHC with mouse samples at this time.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	Lot specific		
Buffer:	PBS, 0.09 % sodium azide		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Storage:	4 °C,-20 °C		
Storage Comment:	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.		
Expiry Date:	6 months		



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Human Spleen (formalin-fixed, paraffinembedded) stained with PIM3 antibody ABIN462148 followed by biotinylated goat anti-rabbit IgG secondary antibody ABIN481713, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry

Image 2. Anti-PIM3 antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody concentration 10 ug/ml.