

Datasheet for ABIN674186 anti-Doublecortin antibody (Biotin)



Overview

Overview	
Quantity:	100 μL
Target:	Doublecortin (DCX)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Doublecortin antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human Doublecortin
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	Doublecortin (DCX)
Alternative Name:	Doublecortin (DCX Products)
Background:	Synonyms: Doublecortex, DBCN, Dbct, DC, Doublin, Lis X, Lissencephalin X, Lissencephaly X linked, Lissencephaly X linked doublecortin, LISX, Neuronal migration protein doublecortin,
	SCLH, XLIS.
	Background: Neuronal Marker Doublecortin (DCX) is a microtubule-associated protein

Target Details

expressed almost exclusively in immature neurons. Neuronal precursors begin to express DCX shortly after exiting the cell cycle, and continue to express DCX for 2-3 weeks as the cells mature into neurons. Downregulation of DCX begins after 2 weeks, and occurs at the same time that these cells begin to express, a marker for mature neurons. Due to the nearly exclusive expression of DCX in developing neurons, this protein has been used increasingly as a marker for neurogenesis. Indeed, the levels of DCX expression increase in response to exercise, which occurs in parallel with increased BrdU labelling, currently a "gold standard" in measuring neurogenesis.

Gene ID:

1641

Application Details

Application Notes:	WB: 1:100-1000, IHC-P: 1:100-500
	Optimal working dilution should be determined by the investigator.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months