antibodies - online.com







anti-XRCC4 antibody (AA 144-193)







_					
U	V	er	VI	е	W

Overview	
Quantity:	100 μL
Target:	XRCC4
Binding Specificity:	AA 144-193
Reactivity:	Human, Cow, Guinea Pig, Horse, Zebrafish (Danio rerio), Bat, Monkey, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This XRCC4 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	

Аррисацоп.	western blotting (wb)	
Product Details		
Immunogen:	Synthetic peptide located between aa144-193 of human XRCC4 (Q13426, NP_071801). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan, Gibbon, Baboon, Monkey, Marmoset, Panda, Bovine, Bat, Horse, Pig, Guinea pig (100%), Galago, Mouse, Rat, Dog, Rabbit (92%), Lizard (85%), Opossum, Zebra finch, Chicken, Platypus, Zebrafish (84%).	
	Type of Immunogen: Synthetic peptide	
Specificity:	Human XRCC4	
Predicted Reactivity:	Percent identity by BLAST analysis: Human, Bovine, Horse, Pig, Guinea pig (100%) Mouse, Rat, Dog, Rabbit (92%) Chicken, Zebrafish (84%).	
Purification:	Immunoaffinity purified	

Target Details

Torquit	VDCC4	
Target:	XRCC4	
Alternative Name:	XRCC4 (XRCC4 Products)	
Background:	Name/Gene ID: XRCC4	
	Family: DNA repair	
	Synonyms: XRCC4, DNA repair protein XRCC4	
Gene ID:	7518	
NCBI Accession:	NP_071801	
UniProt:	Q13426	
Pathways:	DNA Damage Repair, Production of Molecular Mediator of Immune Response	
Application Details		
Application Notes:	Approved: WB (0.2 - 1 μg/mL)	
	Hoose Western Dist. Currented dilution at 1 up/pol in E % aking wills / DDC huffer and HDD	
	Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.	
	conjugated and nassitings and a se anated in 1. co,000 100,000 as decondary unitsody.	
	Not recommended for: IHC-P	
Comment:	Target Species of Antibody: Human	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Distilled water	
Concentration:	Lot specific	
Buffer:	Lyophilized from PBS with 2 % sucrose	
Handling Advice:	Avoid repeat freeze-thaw cycles.	
Storage:	4 °C,-20 °C	
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long	
	term use (up to 1 year)	

Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.

Images




Image 1.