antibodies -online.com







anti-AFF2 antibody (N-Term)



Image



| Overview | | |
|-----------------------|--|--|
| Quantity: | 100 μL | |
| Target: | AFF2 | |
| Binding Specificity: | N-Term | |
| Reactivity: | Human, Cow, Rabbit, Horse, Rat, Pig, Bat, Monkey | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This AFF2 antibody is un-conjugated | |
| Application: | Western Blotting (WB) | |
| Product Details | | |
| Immunogen: | Synthetic peptide from N-Terminus of human AFF2 (P51816, NP_002016). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey, Galago, Bovine, Bat, Rabbit, Horse, Pig (100%), Orangutan, Mouse, Guinea pig (92%), Rat, Elephant (85%), Stickleback (84%). | |
| | Type of Immunogen: Synthetic peptide | |
| Specificity: | Human AFF2 | |
| Predicted Reactivity: | Percent identity by BLAST analysis: Human, Bovine, Rabbit, Horse, Pig (100%) Mouse, Guinea pig (92%) Rat (85%). | |
| Purification: | Immunoaffinity purified | |

Target Details

| Target: | AFF2 |
|-------------------|--|
| Alternative Name: | AFF2 / OX19 (AFF2 Products) |
| Background: | Name/Gene ID: AFF2 |
| | Synonyms: AFF2, AF4/FMR2 family, member 2, AF4/FMR2 family member 2, FMR2, Fragile X mental retardation 2, FRAXE, MRX2, Protein 0x19, FMR2P, 0X19, Protein FMR-2 |
| Gene ID: | 2334 |
| NCBI Accession: | NP_002016 |
| UniProt: | P51816 |

Application Details

| Application Notes: | Approved: WB (0.2 - 1 μg/mL) | |
|--------------------|---|--|
| | Usage: Western Blot: Suggested dilution at 1 μ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody. | |
| Comment: | Target Species of Antibody: Human | |
| Restrictions: | For Research Use only | |
| Handling | | |

| Format: | Lyophilized |
|------------------|--|
| Reconstitution: | Distilled water |
| Concentration: | Lot specific |
| Buffer: | Lyophilized from PBS with 2 % sucrose |
| Handling Advice: | Avoid repeat freeze-thaw cycles. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long |
| | term use (up to 1 year) |
| | Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles. |




Image 1.