



[Go to Product page](#)

Datasheet for ABIN6743179
anti-TRIML1 antibody (AA 360-409)

1 Image

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | TRIML1 |
| Binding Specificity: | AA 360-409 |
| Reactivity: | Human, Mouse, Rabbit, Cow, Rat, Dog, Guinea Pig, Monkey, Pig, Bat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TRIML1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | Synthetic peptide located between aa360-409 of human TRIML1 (Q8N9V2, NP_848651). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey, Galago, Mouse, Rat, Elephant, Panda, Dog, Bovine, Bat, Rabbit, Pig, Guinea pig (100%), Horse (92%). Type of Immunogen: Synthetic peptide |
| Specificity: | Human TRIML1 |
| Predicted Reactivity: | Percent identity by BLAST analysis: Mouse, Rat, Dog, Bovine, Rabbit, Pig, Guinea pig (100%) Horse (92%). |
| Purification: | Immunoaffinity purified |

Target Details

| | |
|-------------------|---|
| Target: | TRIML1 |
| Alternative Name: | TRIML1 (TRIML1 Products) |
| Background: | Name/Gene ID: TRIML1 Synonyms: TRIML1, RNF209, Tripartite motif family-like 1, RING finger protein 209 |
| Gene ID: | 339976 |
| NCBI Accession: | NP_848651 |
| UniProt: | Q8N9V2 |

Application Details

| | |
|--------------------|--|
| Application Notes: | Approved: WB (0.2 - 1 µg/mL) Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody. |
| Comment: | Target Species of Antibody: Human |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---|
| Format: | Lyophilized |
| Reconstitution: | Distilled water |
| Concentration: | Lot specific |
| Buffer: | Lyophilized from PBS with 2 % sucrose |
| Handling Advice: | Avoid repeat freeze-thaw cycles. |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year) Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles. |



Image 1.