antibodies - online.com







anti-CST8 antibody (AA 35-84)



Image



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	CST8
Binding Specificity:	AA 35-84
Reactivity:	Human, Mouse, Rat, Horse, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CST8 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Synthetic peptide located between aa35-84 of human CST8 (060676, NP_005483). Percent
	identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Mouse, Horse (100%),
	Monkey, Marmoset, Dog, Bat (92%), Pig (91%), Bovine (90%), Rabbit (85%), Rat, Elephant (84%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human CST8
Predicted Reactivity:	Percent identity by BLAST analysis: Human, Mouse, Horse (100%) Dog (92%) Pig (91%) Bovine
	(90%) Rabbit (85%) Rat (84%).

Target Details

Target:	CST8
Alternative Name:	CST8 / CRES (CST8 Products)
Background:	Name/Gene ID: CST8
	Synonyms: CST8, CRES, Cystatin-8, CTES5
Gene ID:	Synonyms: CST8, CRES, Cystatin-8, CTES5 10047
Gene ID: NCBI Accession:	

Application Details

Application Notes:	Approved: WB (0.2 - 1 μg/mL)
	Usage: Western Blot: Suggested dilution at 1 μ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year)
	Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.




Image 1.