# antibodies - online.com







## anti-PACRG antibody (AA 216-265)

 $100 \, \mu L$ 



Image



$\sim$			
	N/6	1//r	$I \cap V$

Quantity:

Quay.	·	
Target:	PACRG	
Binding Specificity:	AA 216-265	
Reactivity:	Human, Mouse, Rat, Chicken, Zebrafish (Danio rerio), Cow, Horse, Monkey, Xenopus laevis	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PACRG antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	Synthetic peptide located between aa216-265 of human PACRG (Q96M98, NP_689623).	
	Percent identity by BLAST analysis: Human, Gorilla, Monkey, Galago, Mouse, Rat, Elephant,	
	Panda, Bovine, Horse, Chicken, Xenopus, Salmon, Zebrafish (100%), Opossum, Guinea pig,	
	Panda, Bovine, Horse, Chicken, Xenopus, Salmon, Zebrafish (100%), Opossum, Guinea pig, Zebra finch (92%).	
Specificity:	Zebra finch (92%).	
Specificity: Predicted Reactivity:	Zebra finch (92%).  Type of Immunogen: Synthetic peptide	
	Zebra finch (92%).  Type of Immunogen: Synthetic peptide  Human PACRG	
	Zebra finch (92%).  Type of Immunogen: Synthetic peptide  Human PACRG  Percent identity by BLAST analysis: Human, Mouse, Rat, Bovine, Chicken, Zebrafish (100%)	

#### **Target Details**

Target:	PACRG
Alternative Name:	GLUP / PACRG (PACRG Products)
Background:	Name/Gene ID: PACRG
	Synonyms: PACRG, GLUP, HAK005771, PARK2 co-regulated, PARK2CRG, RP3-495010.2, PARK2 coregulated gene protein
Gene ID:	135138
NCBI Accession:	NP_689623
UniProt:	Q96M98

#### **Application Details**

Application Notes:	Approved: WB (0.2 - 1 μg/mL)
	Usage: Western Blot: Suggested dilution at 1 $\mu$ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

### Handling

Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year)  Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.




Image 1.