antibodies - online.com







anti-Fc epsilon RI/FCER1A antibody (AA 35-84)



Image



\sim	
()\/\	rview
\circ	

Quantity:	100 μL
Target:	Fc epsilon RI/FCER1A (FCER1A)
Binding Specificity:	AA 35-84
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fc epsilon RI/FCER1A antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Jagot Detailo	
Immunogen:	Synthetic peptide located between aa35-84 of human FCER1A (P12319, NP_001992). Percent
	Synthetic peptide located between aa35-84 of human FCER1A (P12319, NP_001992). Percent identity by BLAST analysis: Human, Chimpanzee, Gibbon (100%), Gorilla, Monkey (85%).
	identity by BLAST analysis: Human, Chimpanzee, Gibbon (100%), Gorilla, Monkey (85%).
Immunogen:	identity by BLAST analysis: Human, Chimpanzee, Gibbon (100%), Gorilla, Monkey (85%). Type of Immunogen: Synthetic peptide
Immunogen: Specificity:	identity by BLAST analysis: Human, Chimpanzee, Gibbon (100%), Gorilla, Monkey (85%). Type of Immunogen: Synthetic peptide Human FCER1A
Immunogen: Specificity: Predicted Reactivity:	identity by BLAST analysis: Human, Chimpanzee, Gibbon (100%), Gorilla, Monkey (85%). Type of Immunogen: Synthetic peptide Human FCER1A Percent identity by BLAST analysis: Human (100%).
Immunogen: Specificity: Predicted Reactivity: Purification:	identity by BLAST analysis: Human, Chimpanzee, Gibbon (100%), Gorilla, Monkey (85%). Type of Immunogen: Synthetic peptide Human FCER1A Percent identity by BLAST analysis: Human (100%).

Target Details

Background:	Name/Gene ID: FCER1A
	Family: Immunoglobulin
	Synonyms: FCER1A, Alpha polypeptide, FCE1A, FcERI, Fc Epsilon RI, Fc epsilon RI alpha-chain
	Fc-epsilon RI-alpha, IgE Fc receptor subunit alpha
Gene ID:	2205
NCBI Accession:	NP_001992
UniProt:	P12319
Pathways:	Fc-epsilon Receptor Signaling Pathway, Regulation of Leukocyte Mediated Immunity, Positive
	Regulation of Immune Effector Process
Application Details	
Application Notes:	Approved: WB (0.2 - 1 μg/mL)
	Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP
	conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long
	term use (up to 1 year)
	Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.




Image 1.