

Datasheet for ABIN6746040

## anti-Interleukin enhancer-binding factor 3 (ILF3) (AA 731-780) antibody



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	100 µL
Target:	Interleukin enhancer-binding factor 3 (ILF3)
Binding Specificity:	AA 731-780
Reactivity:	Human, Mouse, Dog, Guinea Pig, Pig, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	Western Blotting (WB)

#### Product Details

Immunogen:	Synthetic peptide located between aa731-780 of mouse Ilf3 (Q9Z1X4, NP_034691). Percent identity by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Mouse, Elephant, Dog, Pig, Opossum, Guinea pig (100%), Rat, Bovine, Rabbit, Horse (92%), Bat (84%).  Type of Immunogen: Synthetic peptide
Specificity:	Mouse ILF3
Predicted Reactivity:	Percent identity by BLAST analysis: Human, Dog, Pig, Guinea pig (100%) Mouse, Rat, Bovine, Rabbit, Horse (92%).
Purification:	Immunoaffinity purified

## Target Details

---

Target:	Interleukin enhancer-binding factor 3 (ILF3)
Alternative Name:	NF90 / ILF3 ( <a href="#">ILF3 Products</a> )
Background:	Name/Gene ID: ILF3  Synonyms: ILF3, CBTF, DRBF, DRBP76, MMP4, NF-AT-90, NF90, NFAR2, Nuclear factor nf90, M-phase phosphoprotein 4, TCP110, MPHOSPH4, NF110b, NF90a, NF90b, NFAR-1, TCP80, NF110, NFAR
Gene ID:	3609
NCBI Accession:	<a href="#">NP_034691</a>
UniProt:	<a href="#">Q12906</a>
Pathways:	<a href="#">M Phase</a>

## Application Details

---

Application Notes:	Approved: WB (0.2 - 1 µg/mL)  Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.
Comment:	Target Species of Antibody: Mouse
Restrictions:	For Research Use only

## Handling

---

Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year)  Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.



**Image 1.**