antibodies .- online.com





anti-DUSP11 antibody (AA 41-90)



Image

Publications



U	V	er	V	ie	W

Quantity:	100 μL
Target:	DUSP11
Binding Specificity:	AA 41-90
Reactivity:	Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DUSP11 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Synthetic peptide located between aa41-90 of mouse Dusp11 (Q6NXK5, NP_082375). Percent
	identity by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Rabbit (100%),
	Elephant, Bovine, Bat, Horse, Pig (92%), Galago, Dog, Opossum (85%), Guinea pig (84%).
	Type of Immunogen: Synthetic peptide
Specificity:	Mouse DUSP11
Predicted Reactivity:	Percent identity by BLAST analysis: Human, Rabbit (100%) Rat, Horse, Pig (92%) Dog (85%) Guinea pig (84%).
Purification:	

Target Details

Target:	DUSP11
Alternative Name:	DUSP11 (DUSP11 Products)
Background:	Name/Gene ID: DUSP11 Subfamily: Protein Phosphatase - Dual specificity other
	Synonyms: DUSP11, PIR1, Dusp 11
Gene ID:	Synonyms: DUSP11, PIR1, Dusp 11 8446
Gene ID: NCBI Accession:	

Application Details

Application Notes:	Approved: WB (0.2 - 1 μg/mL)
	Usage: Western Blot: Suggested dilution at 1 μ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.
Comment:	Target Species of Antibody: Mouse
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year) Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.

Product cited in:

Shahan, Sorenson, Simpson, Kefalides, Lewis: "Tyrosine kinase activation in response to fungal spores is primarily dependent on endogenous reactive oxygen production in macrophages." in: **The Journal of biological chemistry**, Vol. 275, Issue 14, pp. 10175-81, (2000) (PubMed).

Yang, Malek, Desiderio: "An SH3-binding site conserved in Bruton's tyrosine kinase and related tyrosine kinases mediates specific protein interactions in vitro and in vivo." in: **The Journal of biological chemistry**, Vol. 270, Issue 35, pp. 20832-40, (1995) (PubMed).

Aoki, Isselbacher, Pillai: "Bruton tyrosine kinase is tyrosine phosphorylated and activated in pre-B lymphocytes and receptor-ligated B cells." in: **Proceedings of the National Academy of Sciences of the United States of America**, Vol. 91, Issue 22, pp. 10606-9, (1994) (PubMed).

Vetrie, Vorechovský, Sideras, Holland, Davies, Flinter, Hammarström, Kinnon, Levinsky, Bobrow: "The gene involved in X-linked agammaglobulinaemia is a member of the src family of proteintyrosine kinases." in: **Nature**, Vol. 361, Issue 6409, pp. 226-33, (1993) (PubMed).

Images




Image 1.