

Datasheet for ABIN6746874
anti-TBRG1 antibody (N-Term)



[Go to Product page](#)

1 Image **3** Publications

Overview

Quantity:	100 µL
Target:	TBRG1
Binding Specificity:	N-Term
Reactivity:	Human, Horse, Rabbit, Pig, Cow, Dog, Guinea Pig, Monkey, Hamster
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBRG1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Synthetic peptide from N-Terminus of human TBRG1 (Q3YBR2, NP_116200). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey, Galago, Marmoset, Hamster, Panda, Dog, Bovine, Rabbit, Horse, Pig, Opossum, Guinea pig (100%), Mouse, Rat, Elephant (92%), Bat (85%). Type of Immunogen: Synthetic peptide
Specificity:	Human TBRG1
Predicted Reactivity:	Percent identity by BLAST analysis: Human, Dog, Bovine, Horse, Pig (100%) Mouse, Rat (92%).
Purification:	Immunoaffinity purified

Target Details

Target:	TBRG1
Alternative Name:	TBRG1 (TBRG1 Products)
Background:	Name/Gene ID: TBRG1 Synonyms: TBRG1, NIAM, TB-5
Gene ID:	84897
NCBI Accession:	NP_116200
UniProt:	Q3YBR2

Application Details

Application Notes:	Approved: WB (0.2 - 1 µg/mL) Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year) Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.

Publications

Product cited in:	Shain, Landowski, Dalton et al.: "Adhesion-mediated intracellular redistribution of c-Fas-
-------------------	--

Publications

associated death domain-like IL-1-converting enzyme-like inhibitory protein-long confers resistance to CD95-induced apoptosis in hematopoietic ..." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 168, Issue 5, pp. 2544-53, (2002) ([PubMed](#)).

Brown, Holden, Rahn, Perkins: "Immunohistochemical staining for DNA topoisomerase IIa in Hodgkin's disease." in: **American journal of clinical pathology**, Vol. 109, Issue 1, pp. 39-44, (1998) ([PubMed](#)).

Herzog, Holmes, Tuschong, Ganapathi, Zwelling: "Absence of topoisomerase IIbeta in an amacrine-resistant human leukemia cell line with mutant topoisomerase IIalpha." in: **Cancer research**, Vol. 58, Issue 23, pp. 5298-300, (1998) ([PubMed](#)).

Images




Image 1.