antibodies -online.com







anti-RNF139 antibody (AA 72-121)





\sim			
	N/6	1//r	$I \cap V$

Quantity:	100 μL
Target:	RNF139
Binding Specificity:	AA 72-121
Reactivity:	Human, Mouse, Rat, Dog, Rabbit, Cow, Guinea Pig, Bat, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF139 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic peptide located between aa72-121 of human RNF139 (Q8WU17, NP_009149). Percent identity by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Monkey, Galago, Mouse, Rat, Elephant, Dog, Bovine, Bat, Pig, Opossum, Guinea pig (100%), Marmoset, Horse, Platypus (92%), Rabbit (86%).
	Type of Immunogen: Synthetic peptide
Isotype:	IgG
Specificity:	Human RNF139

Product Details (86%). Purification: Immunoaffinity purified **Target Details RNF139** Target: Alternative Name: RNF139 / TRC8 (RNF139 Products) Background: Name/Gene ID: RNF139 Synonyms: RNF139, HRCA1, RCA1, TRC8, Ring finger protein 139 Gene ID: 11236 NCBI Accession: NP_009149 UniProt: **Q8WU17 Application Details** Application Notes: Approved: IHC, IHC-P (5 - 10 µg/mL), WB Usage: ELISA titer using peptide based assay: 1:312500. Western Blot: Suggested dilution at 1 μ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1:50000 - 100000 as second antibody. Comment: Target Species of Antibody: Human Restrictions: For Research Use only Handling

Format:	Lyophilized	
Reconstitution:	Distilled Water.	
Concentration:	Lot specific	
Buffer:	Lyophilized from PBS with 2 % sucrose	
Handling Advice:	Avoid repeat freeze-thaw cycles.	
Storage:	4 °C,-20 °C	
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long	

term use (up to 1 year)

Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.

Images




Image 1.





Image 2.



Immunohistochemistry

Image 3. Anti-RNF139 / TRC8 antibody IHC of human adrenal. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody dilution 5-10 ug/ml. This image was taken for the unconjugated form of this product. Ot ...