# antibodies -online.com









Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μL
Target:	ICK
Binding Specificity:	AA 448-497
Reactivity:	Human, Horse, Rabbit, Dog, Bat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ICK antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

#### **Product Details**

Brand:	IHC-plus™	
Immunogen:	Synthetic peptide located between aa448-497 of human ICK (Q9UPZ9, NP_055735). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Dog, Bat, Rabbit, Horse (100%), Orangutan, Monkey, Marmoset, Elephant, Panda, Bovine, Guinea pig (92%), Galago, Pig, Opossum (85%).	
	Type of Immunogen: Synthetic peptide	
Specificity:	Human ICK	
Predicted Reactivity:	Percent identity by BLAST analysis:	
Purification:	Immunoaffinity purified	

### **Target Details**


Target:	ICK
Alternative Name:	ICK (ICK Products)
Background:	Name/Gene ID: ICK Subfamily: MAK Family: Protein Kinase
	Synonyms: ICK, HICK, Intestinal cell kinase, MAK-related kinase, MRK, Mrk kinase, Laryngeal cancer kinase 2, LCK2, ECO, KIAA0936
Gene ID:	22858
NCBI Accession:	NP_055735
UniProt:	Q9UPZ9

# **Application Details**


Application Notes:	Approved: IHC, IHC-P (5 - 10 μg/mL), WB
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

# Handling

Format:	Lyophilized
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year)
	Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.



#### Image 1.



#### **Immunohistochemistry**

**Image 2.** Anti-ICK antibody IHC of human small intestine. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody dilution 5-10 ug/ml. This image was taken for the unconjugated form of this product. Othe ...