

Datasheet for ABIN675443
anti-PLAUR antibody (AA 201-300)

5 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	PLAUR
Binding Specificity:	AA 201-300
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLAUR antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PLAUR
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep
Purification:	Purified by Protein A.

Target Details

Target:	PLAUR
Alternative Name:	PLAUR (PLAUR Products)

Target Details


Background:	Synonyms: CD87, UPAR, URKR, U-PAR, Urokinase plasminogen activator surface receptor, Monocyte activation antigen Mo3, PLAUR, MO3 Background: Acts as a receptor for urokinase plasminogen activator. Plays a role in localizing and promoting plasmin formation. Mediates the proteolysis-independent signal transduction activation effects of U-PA. It is subject to negative-feedback regulation by U-PA which cleaves it into an inactive form.
Gene ID:	5329
UniProt:	Q03405
Pathways:	Inositol Metabolic Process

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 FCM 1:20-100 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months




Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded rat brain labeled with Anti-PLAUR Polyclonal Antibody, Unconjugated (ABIN675443) at 1:200 followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin-fixed and paraffin embedded mouse brain labeled with Anti-PLAUR Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin embedded mouse adrenal gland labeled with Anti-PLAUR Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN675443.