

Datasheet for ABIN6755413

anti-HMGB1 antibody (N-Term)





Go to Product page

_						
	1//	Д	rv	16	٦/	٨
	W	\vdash	ΙV	Ιt	٦,	/V


Quantity:	100 μL
Target:	HMGB1
- Target.	TIWODI
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog, Pig, Guinea Pig, Horse, Rabbit, Monkey, Chicken, Hamster, Bat,
	Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGB1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-
	embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™	
Immunogen:	Synthetic peptide from N-Terminus of human HMGB1 (P09429, NP_002119). Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Baboon, Monkey, Galago, Marmoset, Mouse, Rat, Shrew, Hamster, Elephant, Dog, Bovine, Bat, Rabbit, Horse, Pig, Guinea pig, Turkey, Zebra finch, Chicken, Platypus, Lizard, Xenopus, Frog (100%), Panda, Opossum (92%).	
	Type of Immunogen: Synthetic peptide	
Specificity:	Human HMGB1	
Purification:	Immunoaffinity purified	

Target Details

Target:	HMGB1	
Alternative Name:	HMG1 / HMGB1 (HMGB1 Products)	
Background:	Name/Gene ID: HMGB1	
	Synonyms: HMGB1, Amphoterin, High mobility group protein B1, High mobility group box 1, High mobility group protein 1, HMG1, HMG3, SBP-1, High-mobility group box 1, HMG-1	
Gene ID:	3146	
NCBI Accession:	NP_002119	
Pathways:	p53 Signaling, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development, Positive Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process, Toll-Like Receptors Cascades, Smooth Muscle Cell Migration, Inflammasome	
Application Details		
Application Notes:	Approved: IHC, IHC-P (10 μg/mL), WB (0.2 - 1 μg/mL)	
	Usage: Western Blot: Suggested dilution at 1 µg/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1: 50,000 - 100,000 as secondary antibody.	
Comment:	Target Species of Antibody: Human	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Distilled water	
Concentration:	Lot specific	
Buffer:	Lyophilized from PBS with 2 % sucrose	
Handling Advice:	Avoid repeat freeze-thaw cycles.	
Storage:	4 °C,-20 °C	
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long term use (up to 1 year) Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.	



Immunohistochemistry

Image 1. Human Colon: Formalin-Fixed, Paraffin-Embedded (FFPE). This image was taken for the unconjugated form of this product. Other forms have not been tested.