# antibodies -online.com





# anti-ZW10 antibody (Biotin)



Go to Product page

$\sim$					
( )	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	ZW10	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZW10 antibody is conjugated to Biotin	
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from mouse ZW10	
Isotype:	IgG	
Predicted Reactivity:	cted Reactivity: Human,Mouse,Rat	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	ZW10
Abstract:	ZW10 Products
Background:	Synonyms: Centromere/kinetochore protein zw10, Centromere/kinetochore protein zw10 homolog, HZW 10, HZW10, Kinetochore associated homolog, KNTC1AP, MGC149821, Zeste
	White 10, Zeste white 10 homolog, ZW 10, ZW 10, centromere/kinetochore protein, ZW 10

kinetochore associated homolog, ZW10 Drosophila homolog centromere/kinetochore protein, ZW10 homolog centromere/kinetochore protein Drosophila, ZW10 homolog centromere/kinetochore protein, ZW10 kinetochore associated homolog, ZW10\_MOUSE. Background: The mitotic checkpoint ensures that chromosomes are divided equally between daughter cells and is a primary mechanism preventing the chromosome instability often seen in aneuploid human tumors. This gene encodes a protein that is one of many involved in mechanisms to ensure proper chromosome segregation during cell division. The encoded protein binds to centromeres during the prophase, metaphase, and early anaphase cell division stages and to kinetochore microtubules during metaphase. It is part of the MIS12 complex, which may be fundamental for kinetochore formation and proper chromosome segregation during mitosis. In mitotic human cells ZW10 resides in a complex with Rod and Zwilch, whereas another ZW10 partner, Zwint-1, is part of a separate complex of structural kinetochore components including Mis12 and Ndc80-Hec1. Zwint-1 is critical for recruiting ZW10 to unattached kinetochores. Depletion from human cells demonstrates that the ZW10 complex is essential for stable binding of a Mad1-Mad2 complex to unattached kinetochores. Thus, ZW10 functions as a linker between the core structural elements of the outer kinetochore and components that catalyze generation of the mitotic checkpoint-derived "stop anaphase" inhibitor.

Gene ID:

9183

#### Application Details

Application Notes: WB 1:100-1000

IHC-P 1:100-500

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	