antibodies -online.com





anti-ABCB11 antibody (AA 21-120)

3 Images



Go to Product page

_					
U	V	er	V	Ie	W

Quantity:	100 μL
Target:	ABCB11
Binding Specificity:	AA 21-120
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ABCB11 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse BSEP
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Rabbit
Purification:	Purified by Protein A.

Target Details


Target:	ABCB11
Alternative Name:	BSEP/ABCB11 (ABCB11 Products)
Background: Synonyms: Bsep, PGY4, SPGP, ABC16, Lith1, PFIC2, Bile salt export pump, ATP-bindir	

Target Details

Expiry Date:


12 months

Target Details	
	sub-family B member 11, Sister of P-glycoprotein, Abcb11 Background: Involved in the ATP-dependent secretion of bile salts into the canaliculus of hepatocytes.
Gene ID:	27413
UniProt:	Q9QY30
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.




Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded rat pancreatic tissue labeled with Anti-BSEP/ABCB11 Polyclonal Antibody, Unconjugated (ABIN675848) at 1:200 followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin-fixed and paraffin embedded rat pancreatic tissue labeled with Anti-BSEP/ABCB11 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Paraformaldehyde-fixed, paraffin embedded human liver carcinoma, Antigen retrieval by boiling in sodium citrate buffer (pH6) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes, Blocking buffer (normal goat serum) at 37°C for 20min, Antibody incubation withRabbit Anti-BSEP Polyclonal Antibody, Unconjugated at 1:500 overnight at 4°C, followed by a conjugated secondary for 90 minutes and DAB staining.