antibodies -online.com





anti-SLC8A2 antibody (AA 601-700) (Cy7)



Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | SLC8A2 |
| Binding Specificity: | AA 601-700 |
| Reactivity: | Rat, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SLC8A2 antibody is conjugated to Cy7 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human SLC8A2 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Predicted Reactivity: | Human,Dog,Cow,Pig,Horse,Rabbit,Guinea Pig |
| Purification: | Purified by Protein A. |

Target Details

| Target: | SLC8A2 |
|-------------------|--------------------------|
| Alternative Name: | Slc8a2 (SLC8A2 Products) |

Target Details

| raiget Details | |
|---------------------|---|
| Background: | Synonyms: NCX2, Sodium/calcium exchanger 2, Na(+)/Ca(2+)-exchange protein 2, Solute carrier family 8 member 2, SLC8A2, KIAA1087 Background: Rapidly transports Ca(2+) during excitation-contraction coupling. Ca(2+) is extruded from the cell during relaxation so as to prevent overloading of intracellular stores (By similarity). |
| Gene ID: | 6543 |
| UniProt: | Q9UPR5 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |