antibodies .- online.com





Datasheet for ABIN678233

anti-ADAR antibody (AA 1001-1226)



Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | ADAR |
| Binding Specificity: | AA 1001-1226 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ADAR antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human ADAR1 |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |
| Target Details | |

l arget Details

| Target: | ADAR |
|-------------------|-----------------------|
| Alternative Name: | ADAR1 (ADAR Products) |

Target Details

| Background: |
|-------------|
|-------------|

Synonyms: 136 kDa double stranded RNA binding protein, Adar 1, ADAR, Adar 1, Adenosine deaminase RNA specic 1, Adenosine deaminase RNA specic, Adenosine deaminase that act on RNA, AV242451, Double stranded RNA specic adenosine deaminase, Double-stranded RNA-specic editase Adar, Drada, Dsh, Dsrad, dsRNA adenosine deaminase, EC 3.5.4.-, G1P1, I 4, I4, I4 protein, Interferon induced protein 4, Interferon inducible protein 4, K88dsrbp, mZaADAR, p136, Pre-mRNA adenosine deaminase, RNA adenosine deaminase 1, RNA-editing deaminase 1, RNA-editing enzyme 1.

Background: ADAR1 converts adenosine to inosine in dsRNA, which destabilizes the dsRNA helix. This activity is important for various functions like site-specific RNA editing of transcripts of the glutamate receptors and modifying viral RNA genomes (which may be responsible for hypermutation of certain negative-stranded viruses, e.g., measles virus). ADAR1 also binds to short interfering RNAs (siRNA) without editing them and suppresses siRNA-mediated RNA interference. This protein is ubiquitously expressed, with the highest levels being found in brain and lung.

Gene ID:

103

Pathways:

Protein targeting to Nucleus

Application Details

| ۸ | -1:: | NI-+ |
|----|-----------|--------|
| ΑD | olication | notes. |

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid | |
|----------------|---|--|
| Concentration: | 1 μg/μL | |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |

Handling

| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
|--------------------|---|
| | handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |