antibodies .- online.com





anti-Olfactomedin 1 antibody (AA 101-200) (FITC)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	Olfactomedin 1 (OLFM1)
Binding Specificity:	AA 101-200
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Olfactomedin 1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Noelin
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	Olfactomedin 1 (OLFM1)
Alternative Name:	Noelin (OLFM1 Products)

Target Details

Background:	Synonyms: AMY, NOE1, OlfA, NOELIN1, Noelin, Neuronal olfactomedin-related ER localized	
	protein, Olfactomedin-1, OLFM1, NOEL1	
	Background: The Olfactomedin family comprises a diverse group of secreted glycoproteins,	
	which includes OLFM1 (Noelin-1), OLFM2 (Noelin-2), OLFM3 (Noelin-3), OLFM4 (Noelin-4), tiarin,	
	pancortin, gliomedin and mycocilin. These proteins are implicated in the development of the	
	nervous system. Specifically, OLFM1 and OLFM2 expression is observed in the neural plate and	
	neural crest, as well as in the cranial ganglia in mouse at E8-10, and later in brain tissue and in	
	the zone of polarizing activity in the limb. Overexpression of OLFM1 causes an excess of neural	
	crest emigrations and prolonged neural crest production. OLFM2 participates in the regulation	
	of the development of the anterior nervous system. An Arg144Gln mutation in OLFM2 has been	
	implicated as a possible cause for open-angle glaucoma (OAG).	
Gene ID:	10439	
UniProt:	Q99784	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	