# antibodies .- online.com







# anti-MTDH antibody (AA 131-230) (Biotin)



( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Quantity:	100 μL	
Target:	MTDH	
Binding Specificity:	AA 131-230	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MTDH antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human AEG-1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig
Purification:	Purified by Protein A.

### **Target Details**

Target:	MTDH
Alternative Name:	LYRIC (MTDH Products)

### **Target Details**

Background:	Synonyms: 3D3, AEG1, AEG-1, LYRIC, LYRIC/3D3, Protein LYRIC, 3D3/LYRIC, Astrocyte elevated		
	gene-1 protein, Lysine-rich CEACAM1 co-isolated protein, Metadherin, Metastasis adhesion		
	protein, MTDH		
	Background: Downregulates SLC1A2/EAAT2 promoter activity when expressed ectopically.		
	Activates the nuclear factor kappa-B (NF-kappa-B) transcription factor. Promotes anchorage-		
	independent growth of immortalized melanocytes and astrocytes which is a key component in		
	tumor cell expansion. Promotes lung metastasis and also has an effect on bone and brain		
	metastasis, possibly by enhancing the seeding of tumor cells to the target organ endothelium.		
	Induces chemoresistance.		
Gene ID:	92140		
UniProt:	Q86UE4		
Pathways:	Cellular Response to Molecule of Bacterial Origin, Cell-Cell Junction Organization		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		
Expiry Date:	12 months		